

Peering Down the Barrel of a Bacteriophage Portal: The Genome Packaging and Release Valve in P22

Jinghua Tang,¹ Gabriel C. Lander,^{2,8} Adam Olia,³ Rui Li,⁴ Sherwood Casjens,⁵ Peter Prevelige, Jr.,⁴ Gino Cingolani,⁶ Timothy S. Baker,^{1,7,*} and John E. Johnson^{1,2,*}

¹Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093-0378, USA

²Department of Molecular Biology, The Scripps Research Institute, La Jolla, CA 92037, USA

³Department of Biological Sciences, Purdue University, W. Lafayette, IN 47907, USA

⁴Department of Microbiology, The University of Alabama, Birmingham, Birmingham, AL 35294, USA

⁵Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT 84112, USA

⁶Department of Biochemistry and Molecular Biology, Thomas Jefferson University, 233 S. 10th Street, Philadelphia, PA 19107, USA

⁷Division of Biology, University of California, San Diego, La Jolla, CA 92093-0378, USA

⁸Present address Life Science Division, Lawrence Berkeley National Lab, 1 Cyclotron Road, Berkeley, CA 94720, USA

*Correspondence: tsb@ucsd.edu (T.S.B.), jackj@scripps.edu (J.E.J.)

DOI 10.1016/j.str.2011.02.010

SUMMARY

The encapsidated genome in all double-strand DNA bacteriophages is packaged to liquid crystalline density through a unique vertex in the procapsid assembly intermediate, which has a portal protein dodecamer in place of five coat protein subunits. The portal orchestrates DNA packaging and exit, through a series of varying interactions with the scaffolding, terminase, and closure proteins. Here, we report an asymmetric cryoEM reconstruction of the entire P22 virion at 7.8 Å resolution. X-ray crystal structure models of the full-length portal and of the portal lacking 123 residues at the C terminus in complex with gene product 4 ($\Delta 123$ portal-gp4) obtained by Olia et al. (2011) were fitted into this reconstruction. The interpreted density map revealed that the 150 Å, coiled-coil, barrel portion of the portal entraps the last DNA to be packaged and suggests a mechanism for head-full DNA signaling and transient stabilization of the genome during addition of closure proteins.

INTRODUCTION

Bacteriophages provide exceptionally tractable model systems to understand the fundamental mechanistic principles by which large, macromolecular complexes assemble and function. Genetic studies of P22 (family *Podoviridae*), one of many dsDNA bacteriophages, provided detailed cause and effect relationships among the eleven viral gene products that participate in particle assembly and the subset of nine gene products resident in the mature virion (Botstein et al., 1973). Mutational analyses firmly established the order in which P22 gene products interact during assembly, the folding and assembly pathway of the capsid protein, and the regulation of dsDNA packaging (King et al., 1973). These resulted in flow charts of gene product relationships, but transmission electron micrographs of negatively stained P22 samples yielded insufficient resolution to correlate the detailed genetics with structure (King et al., 1976). The development and increasingly routine use of cryoelectron microscopy (cryoEM)

and icosahedral image reconstruction methods in the mid 1990s led to more detailed pictures of the capsid (Prasad et al., 1993; Thuman-Commike et al., 1996), and more recently, structures of whole (asymmetric) virions at comparable resolution (17–20 Å) (Chang et al., 2006; Lander et al., 2006). Detailed segmentation of the isolated tail machine at sub nanometer resolution was additionally achieved, and pseudoatomic structures were derived by homology modeling for three viral proteins: portal (gp1:82.6 kDa), gp4 (18.0 kDa), and gp10 (52.5 kDa) (Lander et al., 2009). Models of the tail needle (plug) (gp26:24.6 kDa) (Olia et al., 2007b, 2009) and tailspike (gp9:71.9 kDa) (Steinbacher et al., 1996, 1997) proteins based on their crystal structures closely matched features in that cryoEM density map. One region of density extending from the portal to near the particle center was misinterpreted in the earlier asymmetric reconstruction of P22 (Lander et al., 2006) as corresponding to the three ejection (“pilot”) proteins known to be in the particle (Casjens and King, 1974).


P22 is a meta-stable assembly, poised to inject its genome into *Salmonella* host bacteria. A total of 415 copies of gp5, the major capsid protein (46.7 kDa), assemble into a nearly icosahedral, $T = 7$ laevo, quasi-symmetric head. A 12-fold symmetric dodecamer of gp1 (“portal”) occupies one of the vertices of the icosahedron, forming an axial channel through which the dsDNA genome enters and exits the capsid. The portal also serves as the attachment site for the tail machine, formed by four proteins (gp4, gp9, gp10, and gp26), each of which assumes a different oligomeric state with respective subunit stoichiometries in a 12:18:6:3 ratio.

Here, we report the asymmetric cryo-reconstruction of the infectious P22 virion at 7.8 Å resolution and the icosahedrally symmetrized map at 5.0 Å. Models derived from the crystal structures of full-length portal and the $\Delta 123$ portal-gp4 complex (“core-gp4 complex”) described in Olia et al. (2011) now provide an accurate density assignment for the entire reconstruction and a compelling description of the particle dynamics that accompany dsDNA packaging and tail-machine morphogenesis.

RESULTS AND DISCUSSION

Asymmetric Cryo-Reconstruction of Virion

Three-dimensional (3D) reconstructions of the P22 virion (Figure 1A) and capsid (Figure 1B) were computed, respectively,

**Figure 1. Cryo-Reconstructions of P22**

(A) Asymmetric reconstruction of P22 virion at 7.8 Å resolution, with density map thresholded and segmented to highlight individual components: gp4 (pink), gp5 (color-coded radially from inner (blue) to outer (white) radii), gp9 (dark yellow), gp10 (cyan), and gp26 (yellow). Scale bar is 100 Å.


(B) A stereo view of the icosahedrally symmetrized P22 capsid at 5.0 Å resolution. A modified capsid protein model (blue) nicely fits the reconstructed density.

from 21,645 and 18,602 images of unstained, vitrified virions (see **Experimental Procedures**). The density map of the icosahedrally symmetrized, T = 7 *l*aevo capsid showed numerous long, tubular structures with a right-handed helical twist, characteristic of α -helical polypeptide segments rendered at \sim 5 Å resolution (**Figure 1B**). These correspond to the homologous “spine” helices prevalent in the core of the coat protein of bacteriophage HK97 (Wikoff et al., 2000). The capsid reconstruction provided a means to closely monitor steps taken to produce the final asymmetric virion reconstruction, which was computed without imposing any global or local symmetry (Tang et al., 2010).

The atomic model for the full-length portal (Olia et al., 2011) nicely fit the asymmetric virion density map, with the 150 Å coiled-coil barrel structure redefining the scabbard-like tube of

density (previously interpreted as ejection proteins (Lander et al., 2006)) that extends from the portal core to near the center of the head (**Figure 2A**; the entire portal, including the core and barrel, are colored red). The organization of the bulk dsDNA in the head (**Figure 2A**, DNA is green) closely matched what was previously reported (Chang et al., 2006; Lander et al., 2006), and includes ten distinct, concentric layers, all of which are \sim 22 Å apart (**Figure 2B**). The bulk DNA in our P22 structure appears to be rendered at \sim 15 Å resolution because that is the intrinsic resolution of the DNA within the capsid due to the dynamic variation that occurs between particles, regardless of the resolution of the capsid.

The DNA adopts dramatically different structures in the region immediately surrounding the scabbard and inside it (**Figures 2A**

**Figure 2. P22 Virion Structure**

(A) Cross-section (\sim 60 Å thick) through the center of the P22 virion cryo-reconstruction, segmented to highlight individual components: gp1 (red), gp4 (pink), gp5 (color-coded radially from inner blue to outer white), gp9 (dark yellow), gp10 (cyan), and DNA plus pilot proteins (green). Scale bar is 200 Å.

(B) Radial density plot of the spherically averaged phage reconstruction highlights at least ten distinct layers of dsDNA, separated by about 22 Å, within the capsid shell. A thin, central section slab from a map averaged with 12-fold symmetry about the portal axis, aligned with and at the same magnification as the radial density plot above it and depicted with a radial color-ramp (red-to-blue, low-to-high radii), shows a closeup view of the DNA layers.

(C) Magnified view of a slab of the virion density map, segmented to highlight the portal/gp4 complex (gray mesh) wedged inside the capsid (blue, solid density), the DNA (green, solid density) contained within the portal channel, and the ribbon model of the portal/gp4 crystal structure (gp1 and gp4 in red and magenta, respectively) fitted into the portal/gp4 density. The gp1 crystal structure clearly does not fit the density near the base (proximal end) of the barrel (black arrow) and extends \sim 10 Å above the reconstructed density at the distal end. Scale bar is 50 Å.

(D) Ribbon model of entire gp1 crystal structure (red) fit into the segmented portal density (gray mesh). Closeup view of boxed region appears in stereo in **Figure 4**. Scale bar is 50 Å.




Figure 3. Conformational Changes in Portal Occur after DNA Packaging

(A) Fit of unmodified gp1 core ribbon model into P22 virion cryo-reconstruction (gray mesh) highlights interactions that occur with the packaged DNA (green colored mesh) and capsid subunits (blue-colored mesh and blue model). The most significant changes in portal conformation occur at the N terminus and in the loops highlighted by arrows in (B).

(B) Ribbon model of gp1 core crystal structure (E5 to G602, ramped in color from blue to red) nicely fits the reconstructed density (gray mesh) of the isolated tail machine (Lander et al., 2009). Black arrows point to two loops that change conformation upon interaction with capsid. A density threshold of $\sim 1\sigma$ was used for the contours.

and 2C). Near the center of the head where the distal end of the barrel terminates, the bulk DNA is unconstrained and disordered whereas, inside the barrel, the DNA is confined and follows a linear path. The density ascribed to DNA again changes at the transition between the proximal end of the barrel and the portal core where it becomes considerably wider than expected for a single duplex. Density assigned to DNA extends ~ 20 Å below the portal core and the gp4 ring, ending just inside the gp10 ring. This final ~ 20 Å of DNA may correspond to the portion of the genome that is not fully inserted into the capsid when the gp2/3 terminase complex disengages from the portal. As described below, we suggest that conformational changes in the barrel of gp1 may function as a “valve” to retain the DNA in the capsid until the gp26 tail needle (plug), which is the primary gene product known to keep the DNA within the virion (Lander et al., 2009; Olia et al., 2007b), is inserted into the tail machine. This occurs after gp4 and gp10 have associated with the portal, implying that intermediate stabilization of the packaged DNA is required.

Near the portal-capsid interface, the individual dsDNA strands are hexagonally close-packed (marked by dots in Figure 3A) and an obvious, dense and circular ring of dsDNA abuts tightly against the portal subunits just inside the capsid (Figure 3A, green density marked by dot closest to the center of the panel). Comparison of rigid body fits of the portal X-ray model into cryo-reconstructions of the virion (Figure 3A) and isolated tail (Figure 3B) show that the model best fits the isolated tail. Portions of the portal whose conformations change in apparent response to the presence of gp5 subunits, include two loops (Figure 3B, arrows) and residues near the N terminus, which clearly fit density present in the isolated tail (Figure 3B; E5 is first residue in the X-ray model) but not in the virion map. All residues on the portal subunit close enough to interact with dsDNA in the virion are identified in Table 1.

Portal Barrel Exhibits Large Conformational Changes

The structure of the portal barrel clearly differs between virions and crystals. A rigid body fit of the full-length portal X-ray model into the virion cryo-reconstruction positions the distal end of the barrel ~ 10 Å beyond the scabbard density (Figure 2C). Also, at the base of the scabbard where it meets the portal core, the X-ray model splay radially outward beyond the EM density

(Figure 2C (arrow) and D). An additional local 12-fold symmetry averaging step was carried out to improve the signal-to-noise ratio of the reconstructed barrel density. When the EM map is rendered at a higher density threshold, the twelve helices in the X-ray model are seen to fall out of phase (i.e., rotated $\sim 10^\circ$ clockwise as viewed down the barrel axis) with the corresponding tube-like features in the virion map (Figure 4). All three regions of inconsistency between the X-ray model and cryoEM density are accommodated if the model is twisted and the bundle of helices is compressed slightly inward toward the axis of the barrel near its base (see Movie S1 available online).

Table 1. Residues in the Portal that Contact dsDNA and Residues in Gp4 that Contact the Tail Spike proteins

Portal Residues in Contact with dsDNA	Gp4 Residues in Contact with gp9 Tail Spike Subunit A	Subunit B	Subunit C
LEU 8	PRO 30	THR 5	THR 5
LYS 228	GLN 31	LYS 6	LYS 6
ARG 249	GLN 34	GLY 7	GLY 7
ASP 250	ALA 68	ASP 8	ASP 8
ILE 251	GLU 69	VAL 10	
LYS 252		ARG 11	
ASP 253		LEU 14	
VAL 254		ASP 27	
ASP 256		GLU 29	
ASP 257		PRO 30	
GLN 270		GLN 31	
ARG 275		MET 33	
TYR 277		GLN 34	
LYS 290		ASN 65	
	PRO 66		
	PRO 67		
	ALA 68		
	GLU 69		
	ASP 71		

Each tail spike interacts with three adjacent gp4 subunits (lists of contact residues in the three gp4 subunits). Gp4 subunits A, B, and C correspond, respectively, to the green, red, and magenta subunits in Figure 6.




Figure 4. Remodeling of the gp1 Crystal Structure to Fit the P22 Virion Density Map

(A) Stereo, close-up view of the back half of the proximal part of the portal barrel (gray surface) into which the gp1 crystal structure (red, ribbon model) is fitted. The model fits the portal core density quite well (“core” consists of residues 1–602 and only the topmost portion appears in this view at the bottom of the panel). Above the core, the model deviates significantly from the density map. For example, the helices traverse empty (low density) space and then line up out of register with tubular density features that follow a slewed path and clearly define the wall of the barrel. The adjusted model (green) fits the density map better with regions pointed by two arrows (1) and (2) highlighted in the cross-sections shown in (B) and (C). See also Movie S1.

This concerted rearrangement shifts the distal end of the barrel toward the core of the portal by ~10 Å, positioning the helices in register with the tube-like features near the proximal end of the scabbard (Figure 4). As the head fills during packaging, the mounting pressure exerted by the DNA may induce these rearrangements in the barrel structure, which we posit functions as a valve to retain the DNA until the tail machine assembles.

Interaction of Closure Protein (gp4) with Capsid

The atomic model of the core-gp4 complex (Olia et al., 2011) fit the reconstructed 8 Å density map of the isolated tail machine (Lander et al., 2009) with virtually no need for adjustment, indicating that release of the tail machine from virions has no deleterious effect on the portal core structure and associated proteins. Although SDS page analysis of the proteins in the isolated tail machine indicates that the barrel residues (602–725) of the portal are present, there was no density that corresponds to a fully formed barrel, indicating that, in the process of tail machine release from the virions, the coiled-coil structure is disrupted and does not reform in the tail machine storage conditions *in vitro*.

Fitting the X-ray model of the core-gp4 complex into the 7.8 Å virion density map revealed the presence of an unexpected interaction between the extended C terminus of gp4 with both the portal and capsid (Figure 5). A stretch of about 21 gp4 C-terminal residues (aa 130–150) lies wedged between the capsid and portal. A potential multistep mechanism that explains this observation includes: (1) The packaging complex of the small terminase (gp3: 18.6 kDa), and gp2 (57.6 kDa), the ATPase that provides the driving force, associates with the portal and packages the DNA. (2) A head full of DNA generates the signal that allows gp2 to cut the DNA and terminate packaging (Casjens et al., 1992). The pressure created by the highly condensed DNA in the head is likely responsible for changing the barrel conformation and this in turn closes the valve that retains the DNA, transducing the signal to the terminase proteins. (3) Next, gp4 monomers compete with the terminase complex for its attachment site on the portal and displace it. This induces or permits a final change in portal conformation that allows the C terminus of gp4 to invade the portal-capsid interface and thereby anchor it securely to the particle. The recently reported P22 procapsid structure clearly shows scaffold protein wedged between the capsid protein and the portal (Chen et al., 2011). Comparing the portal position in the procapsid and the virion shows that the portal increases its contact with the capsid shell during maturation (Figure S1). We propose that this portion of the scaffold remains in place during dsDNA packaging, allowing access of the gp4 C terminus to the bottom of the portal. When gp4 binds, the scaffold protein is displaced allowing the final conformational change implied by the position of the gp4-C-terminal polypeptide that is wedged between the capsid and portal.

After 12 gp4 subunits bind to the bottom of the portal, a pre-formed hexamer of gp10 attaches to the gp4 ring (Olia et al., 2007a). The exposed gp4-gp10 interfaces form the sites to which six trimers of the tail spike protein (gp9) attach. Spike trimers are bifunctional, having both receptor-binding activity for O-antigenic repeats of polysaccharides on the salmonella cell surface as well as endoglycosidase activity for cleaving glycosidic bonds after attachment (Baxa et al., 1996). A pseudodoatomic model of the gp9-gp4 interaction (Figure 6) indicates that this symmetry-mismatched interface is rigid (Table 1, columns 2, 3, 4). However, a flexible hinge on gp9 confers significant adaptability of the gp9 trimers in their various roles. The gp26 tail needle (plug), the final component added to the tail assembly, interacts only with gp10 and prevents escape of DNA from the capsid (Strauss and King, 1984).

Biological Role of the Portal Barrel

The realization that the barrel is such a prominent component in the phage structure prompted us to use bioinformatics to search for sequences likely to form such structures. A strong coiled-coil propensity (Lupas et al., 1991) occurs in the C-terminal ~130 residues of all sequenced P22-like podoviruses, and we find that 20 of 25 podovirus portal proteins (chosen to span the diversity within the *Podoviridae*) have similar C-terminal, coiled-coil predictions (Figure 7A shows a sampling of these). In contrast, the barrel domain is not found in ϕ 29-like phage (Simpson et al., 2000) and is not predicted to be present in the T7-like phages. A random sample of 30 portal proteins from long-tailed siphoviruses and myovirus bacteriophages revealed that only two had




Figure 5. Interactions among P22 Proteins gp1, gp4, and gp5

(A) Stereo, side view of the portal density (red surface) and the fitted gp4 crystal structure (magenta). The surface renderings in panels A and B were calculated at a threshold of $\sim 2\sigma$. The C-terminal loop portion of gp4 (amino acids 122–150 highlighted in green) extends up the portal core.

(B) Same as (A) with ribbon model (blue) for the capsid and density mesh for gp4 added to show that the C-terminal loop of gp4 wedges between the portal and capsid subunit interface. Arrows on the left of (B) delineate the slab shown in (C) and (D).

(C) Top-down view of the modeled portal/gp4 complex for the region defined by the two arrows shown in (B).

(D) Same as (C), but in stereo and including the capsid model (blue ribbon). Overlap between the gp4 (green) and gp5 (blue) models clearly suggests that portions of gp4 must undergo significant conformational changes when the tail structure is added during phage assembly. See also Figure S1.


such a prediction (Figure 7A shows one of these, for phage T4). Thus, portal proteins with a C-terminal barrel appear to be preferred by podoviruses and are found less frequently in other tailed phages.

The barrel structure also suggested a series of C-terminal deletion experiments to explore its role(s) in the P22 infection cycle. A mutational study of the P22 barrel was monitored by SDS-PAGE and western blot analysis and showed that these truncated portals were incorporated into phage that stably packaged full-length DNA and the ejection proteins. A naturally occurring

ring deletion of the 48 C-terminal residues (aa 677–725) has no apparent deleterious effect on virus infectivity (Bazinet et al., 1988). Larger deletions, however, starting between residues 650 and 602, all showed ~ 10 -fold decrease in the percentage of phage that were infectious as determined from the ratio of plaque forming units (pfu) to particles (Figures 7B and 7C). Thus, truncated barrels appear defective in delivering DNA to the host. Collectively, the above observations suggest two, potentially related, functions for the portal barrel. First, as suggested by Olia et al. (2011), it may help direct ordered packaging of the genome. Although particles with truncated forms of the barrel still package DNA, this may lead to disordered DNA arrangements that interfere with efficient release. Alternatively, a complete or nearly complete barrel may be required to deliver the DNA and pilot proteins to the bacterium efficiently. We suggest that the barrel supports both functions.

Conclusions

Crystallography of the full-length P22 portal and the barrel-truncated portal in complex with the closure protein gp4 combined with a sub nanometer, asymmetric cryo-reconstruction of P22, dramatically changed the static description of the P22 virion and the dynamic requirements for its assembly. Density previously interpreted as the internal ejection proteins (gp7, gp16, gp20) is now seen to be the coiled-coil, portal barrel (residues 602–725). Remarkably such a dramatic and functionally important extension of the portal was not anticipated and was fully recognized only after the crystal structure of the isolated, full-length portal was determined. The adjustments required to fit the crystal structure of the barrel to the cryoEM density provide a satisfying mechanism for portal-sensing of head-full packaging pressure and transient portal restriction of DNA release during the addition of the closure proteins. Still not known is the role of the barrel during DNA packaging and release. Does it function as a nozzle that gyrates to guide the layering of the DNA into the shells that are strikingly visible in the cryo-reconstruction (Figure 2B)? Does it maintain its extended structure during DNA release? These questions can be addressed through further cryoEM studies. Particles with portals containing truncated barrels display 10-fold reduction in infectivity, yet normal DNA content. Moderate resolution cryoEM studies of these particles will show if the ordering of the DNA is disturbed in these particles. Likewise, particles lacking the gp26 tail-needle (protein plug), spontaneously lose their DNA after packaging (Strauss and King, 1984) and cryo-reconstructions of these particles will determine if the coiled-coil conformation is retained after DNA release. Finally, it is intriguing that a substantial change is likely to occur in the capsid-portal interaction following addition of the gp4 closure protein to the DNA-containing particle. The C-terminal region of gp4 is likely to bind to the portal followed by a change in interaction with the capsid, that traps the C-terminal residues in the final conformation. As discussed above, scaffolding protein may play a role in this final conformational adjustment (Chen et al., 2011). The strong association of 12 gp4 subunits with the portal via their C-terminal regions provides a robust platform for the addition of the other gene products in the tail machine. After more than 50 years, P22 continues to reluctantly reveal its structural strategies and reward those investigating them.



EXPERIMENTAL PROCEDURES

P22 virion isolation, purification, and cryoEM imaging were described previously (Lander et al., 2006). The original set of particle images used by Lander (Lander et al., 2006) were automatically boxed from the raw micrograph data. In this study, we visually screened each micrograph to identify and manually box out only those particles most likely to be intact virions. This provided a complete data set of 21,645 images that were used to compute the 3D reconstructions. The icosahedral reconstruction of P22 was computed using the AUTO3DEM program package (Yan et al., 2007) and a subset of these images (18,602 particles). This also included automatic reboxing of all particle

Figure 6. Stereo View of Interactions between P22 Proteins gp4 and gp9

Portion of the P22 virion density map (gray mesh) with rigid body-fitted models of three gp4 subunits from the dodecamer crystal structure (colored green, red, and magenta) and a gp9 trimer (yellow for the spike head domain and orange for the spike tail domain) for one of the six tail spikes.




Figure 7. The Coiled-Coil Is Observed in Many Podoviridae and Is Important for Function in P22

(A) Plot of the conserved, predicted coiled-coil motif in the C termini of portal proteins for several members of the *Podoviridae* family, as determined by the software COILS (Lupas et al., 1991) with a 21 amino acid scan window. The C termini of the proteins are aligned at the right side of the plot. The phages encoding these portal proteins include: P22, Sf6, CUS-3, and APSE-1, divergent members of the P22-like phages; 933W, F116, BA3, Xfas53, P-SSP7, PaP3, and ϕ Eco32, podoviruses that are all distantly related to one another; and T4, a long-tailed myovirus. Each of these portal proteins shows a strong peak of coiled-coil probability within its C-terminal 150–100 aa. For some, including P22 (in dark red), the probability drops in the terminal 70 aa, even though the crystal structure clearly shows an uninterrupted coiled-coil extending up to within 6 aa of the C terminus (Olia et al., 2011). The reason for this discrepancy between predicted and observed coiled-coil in P22 is not known.

(B and C) C-terminal deletions of portal protein reduce phage assembly and infectivity. Phage particles/ml (B) and plaque forming units/ml (C) produced by trans-complementation of a P22 nonsense portal mutant with full-length and portal proteins truncated from residues 603–725 ($\Delta 602$) and 651–725 ($\Delta 650$) as previously described (Moore and Prevelige, 2002). Fractions from a CsCl gradient that displayed peak infectivity were collected. The number of particles/ml was determined by DNA absorbance and the titer was determined on a permissive host. EM and gel analysis indicated that the particles produced with truncated portal were properly tailed and contained full-length DNA as occurs in wild-type (WT) phage.

Crystal structure fitting and modeling were carried out with program O (Jones et al., 1991) and figures were prepared with the graphics program Chimera (Goddard et al., 2007). The published P22 capsid model (Chen et al., 2011) was modified to fit the density of the icosahedral reconstruction. The portal boundary estimation was guided by both the crystal structure and the position of the scabbard-like tube density previously identified and misinterpreted as ejection proteins. The initial reconstruction of the scabbard-like tube density showed the constriction at the barrel base and the shrinking at the top of the barrel. Only after an additional local averaging step employing the 12-fold symmetry as found in the barrel crystal structure, did the barrel reconstruction reveal the helical rod separation, thus showing the twist of the helices from the crystal structure.

ACCESSION NUMBERS

The 3D density maps of the asymmetric and icosahedral reconstruction of P22 virion have been deposited in the EMDB at EBI with accession codes EMD-5231 (asymmetric) and EMD-5232 (icosahedral).

SUPPLEMENTAL INFORMATION

Supplemental Information includes one figure and one movie and can be found with this article online at doi:10.1016/j.str.2011.02.010.

ACKNOWLEDGMENTS

This work was supported by NIH Grants R01 GM054076 to J.E.J.; R37 GM-033050 and 1S10 RR-020016 to TSB; AI074825 to S.C. and R56 AI076509 to G.C. Image data used in this study were recorded at the National Resource for Automated Molecular Microscopy at TSRI, which is supported by the National Institutes of Health (NIH) through the National Center for Research Resources' P41 program (RR17573).

Received: February 1, 2011

Revised: February 23, 2011

Accepted: February 26, 2011

Published online: March 24, 2011

REFERENCES

- Baxa, U., Steinbacher, S., Miller, S., Weintraub, A., Huber, R., and Seckler, R. (1996). Interactions of phage P22 tails with their cellular receptor, *Salmonella* O-antigen polysaccharide. *Biophys. J.* 71, 2040–2048.
- Bazinet, C., Benbasat, J., King, J., Carazo, J.M., and Carrascosa, J.L. (1988). Purification and organization of the gene 1 portal protein required for phage P22 DNA packaging. *Biochemistry* 27, 1849–1856.
- Botstein, D., Waddell, C.H., and King, J. (1973). Mechanism of head assembly and DNA encapsulation in *Salmonella* phage p22. I. Genes, proteins, structures and DNA maturation. *J. Mol. Biol.* 80, 669–695.
- Casjens, S., and King, J. (1974). P22 morphogenesis. I: catalytic scaffolding protein in capsid assembly. *J. Supramol. Struct.* 2, 202–224.
- Casjens, S., Wyckoff, E., Hayden, M., Sampson, L., Eppler, K., Randall, S., Moreno, E.T., and Serwer, P. (1992). Bacteriophage P22 portal protein is part of the gauge that regulates packing density of intravirion DNA. *J. Mol. Biol.* 224, 1055–1074.
- Chang, J., Weigele, P., King, J., Chiu, W., and Jiang, W. (2006). Cryo-EM asymmetric reconstruction of bacteriophage P22 reveals organization of its DNA packaging and infecting machinery. *Structure* 14, 1073–1082.
- Chen, D.H., Baker, M.L., Hryc, C.F., Dimai, F., Jakana, J., Wu, W., Dougherty, M., Haase-Pettingell, C., Schmid, M.F., Jiang, W., et al. (2011). Structural basis for scaffolding-mediated assembly and maturation of a dsDNA virus. *Proc. Natl. Acad. Sci. USA* 108, 1355–1360.
- Goddard, T.D., Huang, C.C., and Ferrin, T.E. (2007). Visualizing density maps with UCSF Chimera. *J. Struct. Biol.* 157, 281–287.
- Grigorieff, N. (2007). FREALIGN: high-resolution refinement of single particle structures. *J. Struct. Biol.* 157, 117–125.
- Jones, T.A., Zou, J.Y., Cowan, S.W., and Kjeldgaard, M. (1991). Improved methods for building protein models in electron density maps and the location of errors in these models. *Acta Crystallogr. A* 47, 110–119.
- King, J., Lenk, E.V., and Botstein, D. (1973). Mechanism of head assembly and DNA encapsulation in *Salmonella* phage P22. II. Morphogenetic pathway. *J. Mol. Biol.* 80, 697–731.
- King, J., Botstein, D., Casjens, S., Earnshaw, W., Harrison, S., and Lenk, E. (1976). Structure and assembly of the capsid of bacteriophage P22. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 276, 37–49.
- Lander, G.C., Tang, L., Casjens, S.R., Gilcrease, E.B., Prevelige, P., Poliakov, A., Potter, C.S., Carragher, B., and Johnson, J.E. (2006). The structure of an infectious P22 virion shows the signal for headful DNA packaging. *Science* 312, 1791–1795.
- Lander, G.C., Khayat, R., Li, R., Prevelige, P.E., Potter, C.S., Carragher, B., and Johnson, J.E. (2009). The P22 tail machine at subnanometer resolution reveals the architecture of an infection conduit. *Structure* 17, 789–799.
- Lupas, A., Van Dyke, M., and Stock, J. (1991). Predicting coiled coils from protein sequences. *Science* 252, 1162–1164.
- Moore, S.D., and Prevelige, P.E., Jr. (2002). Bacteriophage p22 portal vertex formation in vivo. *J. Mol. Biol.* 315, 975–994.
- Olia, A.S., Bhardwaj, A., Joss, L., Casjens, S., and Cingolani, G. (2007a). Role of gene 10 protein in the hierarchical assembly of the bacteriophage P22 portal vertex structure. *Biochemistry* 46, 8776–8784.
- Olia, A.S., Casjens, S., and Cingolani, G. (2007b). Structure of phage P22 cell envelope-penetrating needle. *Nat. Struct. Mol. Biol.* 14, 1221–1226.
- Olia, A.S., Casjens, S., and Cingolani, G. (2009). Structural plasticity of the phage P22 tail needle gp26 probed with xenon gas. *Protein Sci.* 18, 537–548.
- Olia, A.S., Prevelige, P.E., Jr., Johnson, J.E., and Cingolani, G. (2011). Three-dimensional structure of a viral genome-delivery portal vertex. *Nat. Struct. Mol. Biol.*, (in press).
- Prasad, B.V., Prevelige, P.E., Marietta, E., Chen, R.O., Thomas, D., King, J., and Chiu, W. (1993). Three-dimensional transformation of capsids associated with genome packaging in a bacterial virus. *J. Mol. Biol.* 237, 65–74.
- Simpson, A.A., Tao, Y., Leiman, P.G., Badasso, M.O., He, Y., Jardine, P.J., Olson, N.H., Morais, M.C., Grimes, S., Anderson, D.L., et al. (2000). Structure of the bacteriophage ϕ 29 DNA packaging motor. *Nature* 408, 745–750.
- Steinbacher, S., Baxa, U., Miller, S., Weintraub, A., Seckler, R., and Huber, R. (1996). Crystal structure of phage P22 tailspike protein complexed with *Salmonella* sp. O-antigen receptors. *Proc. Natl. Acad. Sci. USA* 93, 10584–10588.
- Steinbacher, S., Miller, S., Baxa, U., Budisa, N., Weintraub, A., Seckler, R., and Huber, R. (1997). Phage P22 tailspike protein: crystal structure of the head-binding domain at 2.3 Å, fully refined structure of the endorhamnosidase at 1.56 Å resolution, and the molecular basis of O-antigen recognition and cleavage. *J. Mol. Biol.* 267, 865–880.
- Strauss, H., and King, J. (1984). Steps in the stabilization of newly packaged DNA during phage P22 morphogenesis. *J. Mol. Biol.* 172, 523–543.
- Tang, J., Olson, N., Jardine, P.J., Grimes, S., Anderson, D.L., and Baker, T.S. (2008). DNA poised for release in bacteriophage ϕ 29. *Structure* 16, 935–943.
- Tang, J., Sinkovits, R.S., and Baker, T.S. (2010). Three-dimensional asymmetric reconstruction of tailed bacteriophage. *Methods Enzymol.* 482, 209–234.
- Thuman-Commike, P.A., Greene, B., Jakana, J., Prasad, B.V., King, J., Prevelige, P.E., Jr., and Chiu, W. (1996). Three-dimensional structure of scaffolding-containing phage p22 procapsids by electron cryo-microscopy. *J. Mol. Biol.* 260, 85–98.
- van Heel, M., and Schatz, M. (2005). Fourier shell correlation threshold criteria. *J. Struct. Biol.* 151, 250–262.
- Wikoff, W.R., Lijas, L., Duda, R.L., Tsuruta, H., Hendrix, R.W., and Johnson, J.E. (2000). Topologically linked protein rings in the bacteriophage HK97 capsid. *Science* 289, 2129–2133.
- Yan, X., Sinkovits, R.S., and Baker, T.S. (2007). AUTO3DEM—an automated and high throughput program for image reconstruction of icosahedral particles. *J. Struct. Biol.* 157, 73–82.