



Isolation of yeast complex IV in native lipid nanodiscs



Irina A. Smirnova^{a,d}, Dan Sjöstrand^a, Fei Li^{b,1}, Markus Björck^a, Jacob Schäfer^a, Henrik Östbye^a, Martin Högbom^{a,e}, Christoph von Ballmoos^c, Gabriel C. Lander^b, Pia Ädelroth^a, Peter Brzezinski^{a,*}

^a Department of Biochemistry and Biophysics, The Arrhenius Laboratories for Natural Sciences, Stockholm University, SE-106 91 Stockholm, Sweden

^b Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, California 92037, USA

^c Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, 3012 Bern, Switzerland

^d Belozersky Institute, Moscow State University, Leninskie Gory 1, Bldg. 40, 119991 Moscow, Russian Federation

^e Department of Chemistry, Stanford University, Stanford, California 94305, United States

ARTICLE INFO

Article history:

Received 17 June 2016

Received in revised form 23 August 2016

Accepted 7 September 2016

Available online 13 September 2016

Keywords:

Bioenergetics

Proton transfer

Membrane protein

Energy conservation

ABSTRACT

We used the amphipathic styrene maleic acid (SMA) co-polymer to extract cytochrome *c* oxidase (Cyt_cO) in its native lipid environment from *S. cerevisiae* mitochondria. Native nanodiscs containing one Cyt_cO per disc were purified using affinity chromatography. The longest cross-sections of the native nanodiscs were 11 nm × 14 nm. Based on this size we estimated that each Cyt_cO was surrounded by ~100 phospholipids. The native nanodiscs contained the same major phospholipids as those found in the mitochondrial inner membrane. Even though Cyt_cO forms a supercomplex with cytochrome *bc*₁ in the mitochondrial membrane, cyt. *bc*₁ was not found in the native nanodiscs. Yet, the loosely-bound Respiratory SuperComplex factors were found to associate with the isolated Cyt_cO. The native nanodiscs displayed an O₂-reduction activity of ~130 electrons Cyt_cO⁻¹ s⁻¹ and the kinetics of the reaction of the fully reduced Cyt_cO with O₂ was essentially the same as that observed with Cyt_cO in mitochondrial membranes. The kinetics of CO-ligand binding to the Cyt_cO catalytic site was similar in the native nanodiscs and the mitochondrial membranes. We also found that excess SMA reversibly inhibited the catalytic activity of the mitochondrial Cyt_cO, presumably by interfering with cyt. *c* binding. These data point to the importance of removing excess SMA after extraction of the membrane protein. Taken together, our data shows the high potential of using SMA-extracted Cyt_cO for functional and structural studies.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Functional studies of membrane proteins typically require isolation of the protein of interest at a sufficiently high concentration. The most widely used approach is solubilization in detergents. In most cases it is possible to find a detergent that fulfils the criteria for a sufficiently pure, structurally intact and, when relevant, active protein. However, different detergents may be optimal for different experimental methods and changes in function may be introduced during the removal of

the native lipids. In addition, small and/or loosely bound protein subunits (components) may dissociate along with the native lipids. To restore more native-like conditions, amphipathic polymers (known as “amphipols”) may be used to replace the detergent [1]. An even more native-like environment can be re-created by reconstitution of the detergent-purified protein into lipid vesicles [2] or in lipid nanodiscs using a surrounding membrane scaffolding protein [3], which allows functional studies under near-native conditions. However, all these methods involve isolation of the membrane protein of interest using detergent as an intermediate step. Recently, a new technique has emerged that allows extraction and isolation of the membrane protein of interest together with a disc of native lipids [4] using amphipathic styrene maleic acid (SMA) co-polymer (for a recent review, see [5]) without the use of any detergent. This approach has been used to isolate a number of different membrane-bound proteins [6–18], including proton pumps, photosynthetic systems, and ion channels. The procedure has also been used to isolate a secondary transporter from *E. coli* for studies using negative stain electron microscopy [19]. The isolated protein-lipid native nanodiscs (we use here the nomenclature suggested by Dörr et al. [5]) are surrounded by a ring of the SMA co-polymer with a thickness of ~1 nm [12]. The total diameter of the native nanodiscs

Abbreviations: Bis-Tris, 2,2-Bis(hydroxymethyl)-2,2',2''-nitrilotriethanol; cyt. *c*, cytochrome *c*; Cyt_cO, cytochrome *c* oxidase; DDM, *n*-dodecyl β-D-maltoside; EDTA, ethylenediaminetetraacetic acid; HEPES, 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid; MES, 2-(*N*-Morpholino)ethanesulfonic acid; PMS, phenazine methosulfate; SDS, sodium dodecyl sulfate; SMA, poly(styrene-*alt*-maleic acid); TMPD, *N,N,N',N'*-tetramethyl-*p*-phenylenediamine; Tris, Tris(hydroxymethyl)aminomethane.

* Corresponding author.

E-mail address: peterb@dbb.su.se (P. Brzezinski).

¹ Current address: Department of Neurology, UCSF School of Medicine, San Francisco, CA 94143, USA; Department of Biochemistry and Biophysics, University of California, San Francisco, CA 94158, USA.

has been found in the range of 10–24 nm, indicating that there is a large degree of variability in the size of the isolated systems and the lipid–protein ratio. In one recent study [8], the SMA co-polymer was shown to disrupt the *Saccharomyces* (*S.*) *cerevisiae* inner mitochondrial membrane, resulting in the formation of native nanodiscs. The extract was shown to display O₂-reduction activity, catalyzed by the Cyt_cO that was present as one of many components in the mixture of all proteins that were extracted from the native inner mitochondrial membrane.

The mitochondrial Cyt_cO is a multisubunit membrane-bound enzyme located in the cristae of the inner mitochondrial membrane. The *S. cerevisiae* Cyt_cO is composed of 11 subunits where subunits I–III form the functionally active core of the enzyme. The Cyt_cO catalyzes oxidation of water-soluble cyt. *c* and reduction of dioxygen to water in the last step of the respiratory chain. Electrons are donated from the more positive (*p*) side, and protons used in the O₂-reduction are taken up from the more negative (*n*) side, resulting in a charge separation across the membrane. Furthermore, the process is linked to proton pumping across the membrane. The *S. cerevisiae* Cyt_cO binds four redox-active co-factors. Electrons from cyt. *c* are first donated to copper A (Cu_A) and are then transferred consecutively to heme *a* and the catalytic site, which consists of heme *a*₃ and copper B (Cu_B). When O₂ reacts with the four electron-reduced Cyt_cO, electrons are transferred in a number of distinct steps forming intermediates that have been identified and characterized in spectroscopic studies (for review, see [20–22]). First, O₂ binds to the reduced heme *a*₃ resulting in formation of state **A**. Next, an electron is transferred from heme *a* to the catalytic site, the O–O bond is broken, and protons are transferred internally within the catalytic site resulting in formation of a ferryl state that, for historical reasons, is called peroxy (**P**). Formation of this **P** state triggers proton uptake from the *n*-side solution to the catalytic site forming the ferryl state, **F**, which is also linked to fractional electron transfer from Cu_A to heme *a*. In the final step of the reaction the electron from the Cu_A-heme *a* equilibrium is transferred to the catalytic site where water is formed, leaving the Cyt_cO in the oxidized state (**O**).

Results from recent studies showed that two additional proteins, called the respiratory supercomplex factors (Rcf) associate with the *S. cerevisiae* Cyt_cO in the native membranes [23–25]. These factors were suggested to facilitate interactions with the cytochrome (cyt.) *bc*₁ complex in the cyt. *bc*₁-Cyt_cO supercomplex. The Rcfs are presumably loosely bound to the Cyt_cO and can also associate with the cyt. *bc*₁ complex. In a recent study we showed that removal of the regulatory Rcf1 protein results in significant changes in the reaction of the reduced Cyt_cO with O₂ [26] and it is therefore important to make sure that this protein co-purifies with the Cyt_cO upon isolation of the enzyme. When using “weak” detergents such as e.g. digitonin, the *S. cerevisiae* Cyt_cO typically co-purifies with the cyt. *bc*₁ complex in a supercomplex (see above). This supercomplex can be dissociated using detergents such as DDM, but use of this detergent may also result in dissociation of weakly bound components (subunits) such as the Rcf1 polypeptide. Consequently, future studies of the purified Cyt_cO would benefit from development of new methods to isolate the *S. cerevisiae* Cyt_cO along with all its regulatory factors, but without cyt. *bc*₁.

Here, we used the SMA co-polymer to isolate native nanodiscs containing a single Cyt_cO from *S. cerevisiae* in each disc. The isolated Cyt_cO contained both Rcf1 and Rcf2, was fully active, and the kinetics of ligand binding and reaction of the reduced Cyt_cO with O₂ were similar to those observed with intact membranes. Another advantage of the new approach to isolate Cyt_cO is that the native nanodiscs contained all the major lipid components typically found in the yeast inner mitochondrial membrane. Consequently, the method allows for investigation of Cyt_cO function in an environment that is similar to that in the native membranes. Our data also show that excess SMA inhibits the Cyt_cO, which points to the importance of removing excess of the polymer after purification.

2. Results and discussion

2.1. Isolation of the native nanodiscs

The *S. cerevisiae* mitochondria containing His-tagged (7-His on Cox13) Cyt_cO were treated with 2% SMA (styrene:maelic acid at a ratio of 3:1) and the Cyt_cO-containing native nanodiscs were isolated via Ni-NTA affinity chromatography as described in Materials and Methods. As shown in **Fig. S1 A** similar SDS-PAGE patterns were observed for the Cyt_cO-containing native nanodiscs as for the detergent-purified Cyt_cO. The three bands corresponding to molecular weights of 40 kDa, 33 kDa and 22 kDa [27] represent the core subunits I, II and III, respectively. The remaining bands are also very similar for the two samples. The bands above 55 kDa represent contaminations or only partly dissociated Cyt_cO complexes, seen both with the SMA native nanodiscs and with the detergent-purified Cyt_cO. The presence of the loosely-bound Rcf1 and Rcf2 proteins in the Cyt_cO-native nanodiscs was shown using Western blotting of the SDS-PAGE gels followed by immunostaining with specific *anti*-Rcf1 and Rcf2 antibodies, respectively (**Fig. S1B,C**). The bright (light) band on the blot area of **Fig. S1B** indicates the position of Rcf1, which is visualized by chemiluminescence. In the case of Rcf2, some cross-reactivity of the *anti*-Rcf2 antibodies was observed for bands with lower molecular weight, where a band corresponding to Rcf2 was positioned at 25 kDa (**Fig. S1C**).

2.2. Size and shape of the native nanodiscs

Negative stain Electron Microscopy (EM) was used to characterize the size and shape of the SMA-Cyt_cO native nanodiscs (**Fig. 1**). **Fig. 1A** shows that the majority of purified SMA-Cyt_cO native nanodiscs distribute in a monodisperse fashion on the grid, a requirement for accurate analysis of particles by EM classification and reconstruction. A small portion of particles form large aggregates, which might be caused by sporadic adverse effect of the uranyl formate stain, and these were excluded from 2D image analyses. Reference-free 2D classification of the particles shows multiple views of the SMA-Cyt_cO native nanodiscs (**Fig. 1B**) that are similar in size and shape to projections of the crystal structure of Cyt_cO from bovine heart (PDB: 3X2Q) without bound lipids (**Fig. 1C**). As shown in **Fig. 1B** and **C**, the Cyt_cO accounts for the majority of the mass in the SMA-Cyt_cO native nanodiscs, with surrounding lipids contributing to a slight increase in particle diameter relative to the forward projections. A histogram of particle diameters (**Fig. 1D**) shows a bimodal distribution, arising from the disparate dimensions observed for the particles depending on their orientation on the grid. On the basis of the analysis outlined above, we conclude that the native nanodiscs contained one Cyt_cO per disc and that the approximate dimensions of the discs were 11 nm × 14 nm.

Fig. S2 shows a size-exclusion chromatography (SEC) analysis of the SMA-Cyt_cO native nanodiscs. From the position of the major cytochrome-containing peak, the size of the particles was estimated to be in the range equivalent to 600–700 kDa, i.e. a relatively broad band. In order to investigate whether this distribution represents a dynamic equilibrium or a distribution of particles with different shapes/molecular weights, we collected fractions around 400 kDa and around 700 kDa, respectively, and re-run these fractions. In both cases we obtained bands corresponding to either 400 kDa or 700 kDa, which shows that there is a distinct distribution of particles in the SEC that is not distinguishable in the EM analysis.

Next, we compared the molecular weight of the native nanodiscs identified in the SEC experiment to the size obtained from the EM analysis. Assuming an average size of 650 kDa (from the SEC experiment) and an average protein (including the lipids) density of 0.8 Da/Å³, the volume of the isolated particles is ~800 nm³. Assuming spherical particles they would have a diameter of ~12 nm, which approximately corresponds to the particle size as seen using EM.




Fig. 1. Negative stain EM of CytcO-SMA native nanodiscs. (A) Representative negative stain EM image of CytcO-SMA native nanodiscs showing regularly sized and monodisperse complexes. (B) Representative 2D class averages of CytcO-SMA native nanodisc particles. Each class average contains ~100 particles. (C) Projections of crystal structure of bovine CytcO in three different views, scaled to the same pixel size as in panel (B). (D) Distribution of particle sizes obtained for 46,764 CytcO-SMA native nanodiscs particles.

As already mentioned above, the particles observed using EM correspond to native nanodiscs containing one CytcO per disc. Based on a structural model of the *S. cerevisiae* CytcO [28] or the structure of the bovine heart CytcO [29], the dimension of a CytcO monomer is approximately 7 nm × 9 nm when observed perpendicular to the membrane surface. As indicated above, the approximate dimensions of the isolated SMA-CytcO discs were 11 nm × 14 nm, where the smaller of these numbers, 11 nm, is presumably the “height” of the CytcO. The SMA ring around each native nanodisc is ~1 nm [12]. Consequently, assuming that the diameter in the plane of the membrane is ~14 nm, each CytcO is surrounded by a ring of lipids extending 1.5–2.5 nm from the enzyme. The surface area of one DOPC lipid is 70 Å² [30], suggesting that ~100 lipids surround each CytcO molecule in the SMA native discs.

2.3. Spectral characteristics and activity

Fig. 2 shows absorption spectra of the reduced and oxidized (“as purified”) native nanodiscs (Fig. 2A), as well as a reduced minus oxidized difference spectrum (Fig. 2B). The difference spectrum displays features characteristic to CytcO; a peak at 603 nm, attributed primarily to heme *a* (with contribution from heme *a*₃), a peak at 445 nm with contributions from both hemes in their reduced states and a negative peak at 418 nm, attributed to the oxidized hemes. The small peak at ~564 nm is presumably a superposition of local maxima at 560 nm and 568 nm originating from hemes *a* and *a*₃, respectively [31]. We did not observe any peaks characteristic to the cyt. *bc*₁ complex, which shows that this complex did not co-purify with the CytcO.

The steady-state activity of the SMA-extracted CytcO was determined by measuring the O₂-reduction rate in the presence of 10 mM ascorbate, 100 μM TMPD, and 50 μM cyt. *c* at pH 7.4. A value of 130 ± 20 e⁻/s (SD, *n* = 3) was obtained when normalized to the concentration of CytcO (cyt. *aa*₃), based on an absorption coefficient for the reduced-minus-oxidized CytcO at 603 nm of 24 mM⁻¹ cm⁻¹ [31]. For comparison, the CytcO activity in intact *S. cerevisiae* mitochondria was 240 ± 20 e⁻/s (*n* = 6). The activity data are summarized in Table 1. Addition of low concentration of DDM (0.05%), which is a detergent typically used for CytcO purification, enhanced the activity of membrane-bound CytcO, both in the native nanodiscs and in mitochondria, by about a factor of 3 (Table 1). The similarity in the response to addition of detergent to both systems indicates that the CytcO structures were the same in both cases. Furthermore, the data demonstrate that a higher activity does not necessarily imply a “better” preparation. This is

particularly evident from earlier studies of mammalian CytcO, which showed that addition of DDM at high concentration results in loss of regulatory subunits concomitantly with an increase in activity [32].

2.4. Lipid analysis

Fig. 3 shows a thin-layer chromatography analysis of the native nanodiscs (lanes 4 and 5), which demonstrates the presence of the same major phospholipid components as in the original *S. cerevisiae* mitochondrial membranes (lane 3), i.e. PI + PS, PE, PC and CL (for full




Fig. 2. Optical absorption spectra of CytcO-SMA native nanodiscs. (A) Absolute absorption spectra of CytcO-SMA native nanodiscs “as purified” (blue) and after reduction with sodium dithionite (red). (B) A difference absorption spectrum of the reduced sample minus that of “as purified”. The concentration of CytcO was 0.95 μM in 0.2 M NaCl, 20 mM HEPES (pH 7.5).

Table 1
Steady state activity of CytcO-SMA native nanodiscs and of CytcO in the *S. cerevisiae* mitochondrial membrane*.

Sample	Ratio heme aa_3 /protein, nmole/mg [#])	CytcO activity, e ⁻ /s	
		No DDM	+0.05% DDM
CytcO-SMA	0.9 ± 0.4 (n = 4)	130 ± 20 (n = 3)	300 ± 40 (n = 3)
	1.1 ± 0.3 (n = 5)	90 ± 20 (n = 3)	190 ± 40 (n = 3)
Mitochondria	0.08 ± 0.04 (n = 6)	240 ± 20 (n = 6)	700 ± 300 (n = 4)

* For CytcO in SMA native nanodiscs, the heme content was calculated either from the difference spectra “reduced-minus-as purified” or from absolute reduced spectrum (shown in italic, using $\epsilon = 39 \text{ mM}^{-1} \text{ cm}^{-1}$ [31] at 604 nm). Accordingly, two different values are given for the steady-state activity, which is normalized to the CytcO concentration. Because the isolated CytcO was slightly reduced in the “as purified” nanodiscs, the values obtained from the difference spectra yielded an underestimation of the heme concentration.

[#] The ratio of heme aa_3 concentration over the protein amount (see “Materials and Methods” section) is a measure of the CytcO purity. For 100% pure CytcO, with a molecular weight of 200 kDa, the ratio is $\frac{1}{2 \cdot 10^5} \text{ mole/g} = 5 \text{ nmole/mg}$. For detergent-purified CytcO we typically obtained a ratio of ~1. The data show that isolation of the CytcO native nanodiscs resulted in an enrichment of the CytcO by a factor of ~10.

description, see sub-section “Analytical methods” in the Materials and Methods section). Similar results were observed earlier after preparation of other native nanodiscs from *S. cerevisiae* [8], from *E. coli* [4,9] or from *Rhodobacter sphaeroides* [18].

2.5. Ligand binding to the catalytic site

Upon addition of carbon monoxide to the reduced CytcO, the CO molecule binds to heme a_3 . The iron ligand bond is photolabile and can be fully dissociated using a short laser flash. The recombination rate of the heme a_3 -CO complex is sensitive to the structure around the catalytic site heme. Fig. 4A shows absorbance changes at 430 nm and 445 nm (minimum and maximum, respectively, in the CO difference spectrum) after flash-induced dissociation of the CO ligand from the CytcO in SMA native nanodiscs. The recombination reaction displayed components with time constants of $160 \pm 20 \mu\text{s}$ and $6.0 \pm 0.1 \text{ ms}$, respectively. The contribution of the faster of the two components was $30 \pm 10\%$ at 445 nm, as observed previously [26] and shown in Fig. 4A, also with the mitochondrial membranes we observed a small (~10% at 445 nm) rapid component with a time constant of ~100 μs (as well as a major kinetic component with a time constant of

~9 ms). This observation indicates that the altered structural state reflected by the faster CO recombination is present also in the native membranes, although at a lower relative concentration compared to that observed with the CytcO-SMA native nanodiscs. Fig. 4B shows a




Fig. 3. Thin-layer chromatogram of chloroform-methanol extracts of the yeast mitochondria and of CytcO-SMA native nanodiscs. From the left to the right: the 1st and the 2nd lanes are lipid standards (0.04 mg CL, 0.05 mg of each of DOPC, DOPE, PI and PS); the 3rd lane is the extract of the yeast mitochondria (the loaded lipids were extracted from a sample that originally contained ~0.6 mg protein); the 4th and 5th lanes are extracts of two preparations of CytcO-SMA. The loaded lipids (at the arrow) were extracted from a sample that originally contained ~0.2 mg protein. The bands in lane 4 are slightly weaker than those in lane 5, presumably because slightly less material was applied in the former. The bands were visualized by iodine staining. See Materials and methods for more details.




Fig. 4. Absorbance changes after the flash-induced dissociation of CO from fully reduced samples. (A) Absorbance changes as a function of time at 445 nm and 430 nm (above and below the zero (dashed) line, respectively, as indicated in the graph) after flash-induced dissociation of CO (at $t = 0$). The traces represent: CytcO-SMA (blue), yeast mitochondria (grey) and yeast CytcO in DDM (red). Experimental conditions: 0.2 M NaCl, 20 mM HEPES (pH 8.0) for CytcO-SMA; 60 mM sorbitol, 20 mM HEPES (pH 7.4) for yeast mitochondria and 0.15 M KCl, 20 mM HEPES (pH 8.0), 0.05% DDM, 10% glycerol for detergent-solubilized CytcO. The temperature was ~22 °C. All amplitudes are normalized to 1 μM reactive enzyme. (B) Kinetic difference spectra for CytcO-SMA native nanodiscs. Amplitudes of the fast and the slow components of CO-recombination kinetic traces are plotted as a function of wavelength and shown in dark green (squares) and dark blue (circles), respectively. The sum of the amplitudes of these two components at 445 nm has been normalized to 1 μM reacting CytcO. A kinetic difference spectrum (CO-reduced - minus - reduced) of the major component (90%, $\tau \approx 9 \text{ ms}$) in yeast CytcO in native mitochondria is shown for comparison (dark red diamonds). This difference spectrum is normalized to approximately overlap with the slow component (dark blue circles). Time constants and contributions of the kinetic components were determined using the software ProK (global fit).

kinetic difference spectrum of the two components, i.e. the amplitudes of the absorbance changes as a function of wavelength in the ranges of 425–455 nm and 575–625 nm. In the Soret region, both phases displayed very similar difference spectra with coinciding isosbestic points, while in the alpha region the peak around 610 nm of the faster component was red-shifted by a few nanometers as compared to the slower component. These data indicate that the two components reflect CO re-binding to Cyt_cO, however, the two Cyt_cO forms display structural differences around the catalytic site, reflected by different recombination rates and slight spectral differences (c.f. the comparison to the reference kinetic difference spectrum obtained with the yeast mitochondria). In this context it should be noted that the rate of CO recombination may change significantly also upon minor structural changes. For example, the structures of this site are almost identical in both the *Rhodobacter sphaeroides* aa₃ and *Thermus thermophilus* ba₃ Cyt_cO, yet the CO recombination rate is almost a factor of 10 slower with the ba₃ than with the aa₃ Cyt_cO [33–35].

With the detergent-purified Cyt_cO, the main CO-recombination component displayed a time constant of 8 ms. In addition, we observed a rapid component with a time constant of ~140 μs and ~30% of the total absorbance change at 445 nm. In this case, however, the sign of the rapid component at 430 nm was opposite to that observed with the SMA-native nanodiscs and native mitochondrial membranes. Thus, with the detergent-purified Cyt_cO, the rapid component represents CO recombination to heme a₃ in a local structural environment that differs from that in the other two samples (SMA-extracted nanodiscs and native membranes, where the Cyt_cO resides in a membrane). Results from an earlier study with mitochondria showed that a rapid component with a similar time constant of ~100 μs and the same spectral features as in detergent-purified Cyt_cO (absorbance decrease at 430 nm) was observed upon removal of the Rcf1 protein [26]. Also this observation indicates that the 100-μs component observed with the detergent-purified Cyt_cO reflects CO binding to a Cyt_cO that is structurally altered. On the basis of this interpretation we conclude that isolation of Cyt_cO using SMA yields Cyt_cO, which displays CO-ligand binding dynamics that is similar to that observed in the native lipid environment.

2.6. Time-resolved kinetics of the oxidative part of the catalytic cycle

Fig. 5 shows absorbance changes associated with reaction of the reduced Cyt_cO, in SMA native nanodiscs, with O₂. In this experiment,




Fig. 5. Absorbance changes associated with reaction of reduced Cyt_cO with O₂. The reaction was studied with three samples: Cyt_cO in SMA native nanodiscs (blue), in native membranes (grey) and in the detergent DDM (red). The O₂ concentrations after mixing were 0.9 mM, 0.6 mM and 0.6 mM, respectively. For experimental conditions, see Fig. 4. The CO ligand was dissociated by a laser flash at a time ~200 ms after mixing (the flash defines $t = 0$). All amplitudes were normalized to 1 μM reactive enzyme. The rate constants were determined using the software ProK.

the reduced Cyt_cO with CO bound to heme a₃ was mixed with an O₂-containing solution. About 200 ms after mixing, the CO ligand was dissociated by means of a laser flash, which allowed O₂ to bind to heme a₃ and the absorbance changes at 445 nm were monitored over time. The unresolved increase in absorbance at $t \approx 0$ is associated with dissociation of the CO ligand. The following decrease in absorbance is associated with binding of O₂ and stepwise oxidation of the Cyt_cO. The first step after binding of O₂ reflects electron transfer from heme a to the catalytic site, with a time constant of ~20 μs [22], which results in breaking of the O–O bond and formation of the so-called “peroxy” state (P_R). At 445 nm we also observed transfer of the fourth electron, originally residing at Cu_A, to the catalytic site resulting in formation of the oxidized Cyt_cO (O state) with a main component with a time constants of ~0.8 ms (we also observed a small fraction absorbance decrease with a time constant of ~14 ms, see slope in Fig. 5). The main component was slightly slower than that observed with the detergent-purified Cyt_cO (~0.4 ms) [22] or with mitochondria (~0.5 ms).

The total absorbance-change amplitude, as shown in Fig. 5, associated with oxidation of the Cyt_cO, was 83% (0.10) and 92% (0.11) (absorbance change from that at $t = 0^+$ to the final level at ~10⁻² s) with the SMA native nanodiscs and the inner mitochondrial membranes, respectively, of that in detergent-purified Cyt_cO (0.12). The smaller extent of oxidation of the Cyt_cO in SMA native nanodiscs may be an apparent effect that is explained by the method used to normalize the traces. A standard practice is to normalize all traces to the CO-dissociation absorbance change amplitude at $t = 0$ (which was done in this case). This procedure assumes that the Cyt_cO population is uniform. However, as discussed above, we did observe two components in the CO-recombination reaction after addition of SMA, indicating that in the SMA native nanodiscs there is a small fraction Cyt_cO with an altered catalytic site. If the absorbance changes in Fig. 5 would be normalized to the fraction “normal” CO recombination (i.e. equivalent to the amplitude of the 6-ms component) then the absorbance changes associated with oxidation of the Cyt_cO would be more similar in all cases in Fig. 5. A more detailed discussion on the slightly larger amplitude observed with the detergent-purified Cyt_cO than with the mitochondria is beyond the scope of this study and will be discussed elsewhere.

2.7. Inhibition of activity by SMA

A complication when using SMA for purification of Cyt_cO (and probably also other membrane proteins) is that interaction of the co-polymer with the Cyt_cO strongly inhibits the O₂-reduction activity of the enzyme. The inhibition effect was reversible because after addition of SMA the activity was restored upon washing the sample with a buffer without SMA. These results indicate that addition of the SMA co-polymer does not result in altering the structure of the Cyt_cO. The data in Fig. 6A shows the O₂-reduction activity measured for different SMA concentrations added to *S. cerevisiae* mitochondria for two concentrations of cyt. c, 0.05 μM and 50 μM, respectively. For the lower cyt. c concentration, the O₂-reduction activity dropped to 50% of the maximum value at ~0.002% SMA. For the higher cyt. c concentration, we first observed a stimulation of the activity, followed by a drop to 50% of the maximum activity at 0.03–0.04% SMA. This drop in activity at lower SMA concentrations for 0.05 μM than for 50 μM cyt. c indicates that SMA interferes with cyt. c binding to the Cyt_cO. Because the effect was reversible, the excess SMA polymer could, for example, interfere with the electrostatically-controlled docking between cyt. c and Cyt_cO on the *p*-side of the membrane. This conclusion is qualitatively consistent with the increase in activity at low SMA concentrations for the high cyt. c concentration, because optimum activity is observed when the electrostatic interactions are intermediate, i.e. upon increasing the ionic strength (at the lowest ionic strength product dissociation is rate limiting) [36].




Fig. 6. Effect of SMA on the activity of CytcO. (A) Steady-state O₂-reduction activity of the *S. cerevisiae* mitochondria in the presence of SMA. The activity is defined by the oxygen-consumption rate measured with equine heart cyt. *c* at 50 μM (black dots) or 0.05 μM (red squares). The activity is presented as relative values compared to that in the absence of SMA (set as 100%). The 100% activities were 250 e⁻/s and 70 e⁻/s, respectively. The IC₅₀ values for 50 μM and 0.05 μM cyt. *c* were 0.032 10⁻²% and 0.0019 10⁻²%, respectively. The concentration of yeast mitochondria protein was 0.2 mg/ml. For other experimental conditions, see the Materials and Methods section. (B,C) Reaction of fully reduced bovine CytcO with oxygen in the absence (black traces) and in the presence (blue traces) of 0.2% SMA, monitored at 445 nm (B) and 605 nm (C). The O₂ concentration after mixing was 0.9 mM. The buffer contained 0.1 M KCl, 20 mM HEPES (pH 8.0), 0.1 mM EDTA and 0.05% DDM. For other experimental conditions, see Fig. 4. The CO ligand was dissociated by a laser flash at *t* = 0, ~200 ms after mixing. All amplitudes were normalized to 1 μM reactive enzyme. The rate constants were determined using the software ProK.

To test whether or not such an excess SMA co-polymer as used in the activity inhibition measurements interferes with proton uptake from the *n*-side of the membrane, we also investigated the reaction of the reduced mitochondrial CytcO from bovine heart and O₂ (Fig. 6B). This enzyme is well characterized and the kinetic components

associated with proton uptake have been identified in the past [37–39]. In the presence of SMA, the kinetic component having a time constant of ~100 μs (absorbance increase in the black traces in Fig. 6BC in the time range ~200–400 μs), associated with electron transfer from Cu_A to heme *a* during the P_R → F reaction, was absent, which indicates that proton uptake was significantly slowed. However, the time constant of the next transition, F → O, (τ ≈ 1 ms) was about the same in the presence as in the absence of SMA, which indicates that the proton uptake was not slowed beyond ~1 ms. In other words, the slowed proton uptake during P_R → F is not the (only) explanation for the observed SMA effect on the catalytic turnover. Furthermore, the total extent of oxidation was smaller with than without SMA added, which indicates that in the presence of SMA, a small CytcO fraction was not oxidized over the time scale of the experiment (c.f. also data with the SMA native nanodiscs in Fig. 5). Taken together, these data indicate that the dramatically slowed turnover activity seen upon addition of SMA most likely originates from interference with formation of the reduced cyt. *c*-oxidized CytcO complex. In addition, the presence of the SMA polymer presumably does slow proton uptake, however, this effect is not manifested in measurements of the turnover rate.

The activity measured with the isolated yeast SMA-CytcO in this work was 130 e⁻/s, i.e. about 50% of that measured with CytcO in mitochondria. The SMA concentration used for extraction of the CytcO native discs was 2% (note that the total protein content during protein extraction is orders of magnitude higher than that in the experiment shown in Fig. 6), but the sample was then washed in multiple steps; first in the affinity chromatography (without any quantitative estimation of the dilution) and then to further lower than SMA concentration by a factor of 5–10 × 10³. Even though the SMA concentration after washing was negligible, we cannot exclude some remaining specific binding of the SMA co-polymer to the CytcO that would interfere with binding of cyt. *c*.

2.8. Comments on the extraction of CytcO

The use of the SMA co-polymer to extract CytcO offers a new method to isolate pure CytcO with negligible contamination from other heme proteins, yet containing the weakly bound and functionally important Rcf1 and Rcf2 proteins (see Introduction). The possibility to isolate only CytcO along with a significant amount of lipids and with the Rcf proteins is surprising given that in *S. cerevisiae* CytcO forms a supercomplex with cyt. *bc*₁ and that the Rcf proteins are presumably weakly bound [23–25,40–49]. These complexes are typically isolated using weak detergents such as digitonin. Given our findings that the CytcO is extracted with a disc of surrounding lipids, one may expect that the CytcO would be found in the discs together with the cyt. *bc*₁ complex. Indeed, we observed a subpopulation of cyt. *bc*₁ in some preparations, but primarily when the cyt. *bc*₁ to CytcO ratio was increased in some batches of the yeast mitochondrial membranes. It is possible that in the native membrane there is an equilibrium of supercomplexes and free CytcOs, as well as cyt. *bc*₁ complexes. The suggested mechanism by which SMA extracts membrane proteins from the native membranes [4] indicates that the polymer enters the hydrophobic part before extraction. It is possible that this mechanism results in shifting the equilibrium of free complexes versus supercomplexes towards free protein components (i.e. weakening the cyt. *bc*₁-CytcO interaction), before extraction of the CytcO native nanodiscs.

3. Conclusions

We present a detergent-free method to extract CytcO from the *S. cerevisiae* inner mitochondrial membranes into native nanodiscs using a SMA co-polymer. The CytcO native nanodiscs displayed significant O₂-reduction activity. In addition, the ligand-binding kinetics and reaction of the reduced CytcO with O₂ were similar to those measured in the native *S. cerevisiae* mitochondrial membranes. The size of the native nanodiscs indicates that ~100 lipid molecules surround each




Fig. 7. Schematic model of a CytcO-lipid SMA native nanodisc. The CytcO structure is based on a model of yeast CytcO [28] and prepared using the software PyMol [59]. Subunits I and II are shown in orange and green, respectively. The picture illustrates the approximate proportions of the lipid disc and of the protein.

CytcO, which presumably yields a native lipid environment (see model in Fig. 7). The lipid composition in the native nanodiscs was similar to that found in the mitochondrial membrane. The preparation yields a relatively homogeneous size distribution. In future studies this method may prove valuable in functional and structural studies.

4. Materials and methods

4.1. Preparation of the SMA co-polymer

Styrene maleic anhydride resin SMA EF30 (Cray Valley, the ratio of styrene to maleic anhydride was 3:1) was solubilized in water by alkali hydrolysis. The SMA powder (15 g) was stirred in 300 ml of 1 M NaOH overnight, the mixture was then heated to boiling with reflux and a stirred for another 4 h. After complete solubilization of the solid resin the mixture was allowed to cool to room temperature. The mixture was acidified with HCl to reach a pH value about 4. The pellet was collected by centrifugation and washed with water three times (about 400 ml each time). Finally, the pellet was dissolved in water by step-wise addition of 6 M NaOH under stirring; the final pH was about 8 and the final volume about 120 ml. Non-dissolved particles were removed by centrifugation. The obtained solution of hydrolyzed SMA was dialyzed against 50 mM Tris-HCl (pH 8.0). The SMA was quantified by drying a sample to constant weight and the volume of the stock solution was adjusted in order to obtain 5% SMA (w/v) concentration.

4.2. Cultivation of *S. cerevisiae* and preparation of mitochondria and SMA-CytcO native nanodiscs

For the construction of a yeast strain expressing Cox13 with a C-terminal His7Strep tag, the wild-type strain W303a (Mat A ade2-1 his3-11,15 trp1-1 leu2-3,112 ura3-1) was transformed by homologous recombination with a PCR fragment coding for a HIS3 cassette and the sequences allowing to exchange the authentic stop codon of COX13 with a 6 amino acid linker followed by a His7/Strep tag and a new stop codon. The resultant strain (MOY764) was verified by PCR and showed normal respiratory growth. The yeast was grown on YPG medium with vigorous shaking at 30 °C according to Meisinger et al. [50]. Isolation of the crude mitochondrial fraction (for the SMA-treatment) and of highly purified mitochondria (for functional studies) from yeast was performed according to the procedure by Meisinger et al. [50] employing Zymolyase-20 T from *Arthro bacter luteus* (Nacalai Tesque). For time-resolved kinetic experiments (see below) the mitochondria were prepared devoid of their outer membrane according to

[51]. Briefly, mitochondria after running the sucrose gradient [50] were treated with a hypotonic medium containing 60 mM sorbitol, 20 mM HEPES (pH 7.4) and mitoplasts were collected by centrifugation at 10,000 g.

For purification of CytcO-SMA native nanodiscs, crude yeast mitochondria containing CytcO with a 7-His-tag on subunit 13 were incubated with 2% SMA (3:1) at a SMA:protein ratio of 2.5 in 0.2 M NaCl, 50 mM Tris-HCl (pH 8.0) (buffer A) at 4 °C for 4 h with a slow rotation on a rotation wheel. Non-solubilized membranes were removed by centrifugation (17,000 g; 20 min; 4 °C) and the supernatant was collected. After this step, SMA was not present in the buffers and before the affinity chromatography the SMA concentration in the supernatant was lowered by a factor of about three by exchange of the buffer for 0.2 M NaCl, 20 mM HEPES (pH 8.0) (buffer B) using concentrator filters (Amicon Ultra, Ultracel-100 K, Merck Millipore). Then, the solution was loaded on a HisTrap HP column (5 ml, GE Healthcare) by overnight cycling. The column was washed with buffer B supplemented with 5 mM imidazole and the CytcO-SMA native discs were eluted with buffer B, supplemented with 20–50 mM imidazole. Finally, the eluted CytcO-SMA native discs were concentrated and imidazole was removed by exchange to buffer B (dilution by a factor of 5–10 × 10³). The CytcO-SMA native discs were stored at 4 °C.

Pure CytcO from *S. cerevisiae* was prepared as described in [52], except that a TS Series Continuous Flow cell disruptor (Constant System Ltd) was used instead of glass beads. The bovine mitochondrial CytcO used for control experiments was prepared as described in [53].

4.3. Size-exclusion chromatography analysis

The SEC analysis of CytcO-SMA nanodiscs was performed on Superose 6 Increase 10/300 GL (GE Healthcare) linked to the Shimadzu Prominence HPLC system with SIL-20 A Autosampler and SPD-M20 A Detector. Buffer B (see above) was used as a liquid phase. A gel filtration calibration kit, HMW (GE Healthcare) was used for the calibration of the column. The data are shown in Fig. S2.

4.4. Analytical methods

The protein concentration was estimated using a Lowry assay in the presence of SDS.

Optical absorption spectra were recorded using Cary 100 UV-VIS or Cary 400 UV-VIS (Agilent Technologies) spectrophotometer. Sodium dithionite was used as a reducing agent. The concentration of hemes *a* and *a*₃ was estimated from the difference “reduced-minus-oxidized” spectra at 603 nm using an absorption coefficient of 24 mM⁻¹ cm⁻¹, or from the reduced CytcO spectrum at 604 nm using an absorption coefficient of 39 mM⁻¹ cm⁻¹ [31].

SDS-PAGE was performed using pre-cast NuPAGE Bis-Tris 4–12% gels with MES-SDS running buffer. Samples were incubated in the NuPAGE SDS sample buffer with 50 mM DTT for 25 min at 50 °C. Western blotting was run with PVDF transfer membranes (0.45 μm; GE Healthcare) in NuPAGE Bicine transfer buffer. All buffers mentioned above in this paragraph were prepared according to recommendations provided by the manufacturer. The voltage was set according to the manufacturer's recommendations. Page Ruler Plus (pre-stained protein ladder, Thermo Scientific) was used as a standard. Brilliant Blue R250 (Sigma) was used for gel staining. Immunostaining was done with *anti-Rcf1* or *anti-Rcf2* as primary antibodies (gift from H. Dawitz and M. Ott, Stockholm University) and *anti-Rabbit IgG*-Peroxidase antibody produced in goat (Sigma) as secondary antibodies. Chemiluminescence upon oxidation of luminol by hydrogen peroxide in the presence of peroxidase was used for visualization of blots (ECL Prime Western Blotting Detection Reagent, Amersham). Imaging of gels and blots was performed using a Luminescent Image Analyzer LAS-1000 plus (Fujifilm).

Lipid extraction and analysis were performed as follows. A volume of 0.25 ml of SMA-CytcO or of the yeast mitochondria suspension (the amount of starting material for CytcO-SMA and mitochondria was assayed as an amount of protein, which was approximately 2 mg or 4 mg, respectively) was mixed with three volumes (0.75 ml) of chloroform-methanol (1:2) and vortexed. Then, one volume of chloroform and one volume of water were added stepwise with vortexing of the samples after each addition. The samples were chilled on ice and spun at 17,000 g at 4 °C for 7 min. The lipid-containing layer was washed three times with two volumes of cold water. If necessary, the lipid fraction was concentrated by evaporating with a stream of nitrogen. Thin-Layer Chromatography (TLC) of the chloroform-methanol extracts was performed on plates of Silica gel 60 (20 × 20 cm; Merck) with the mixture chloroform/methanol/acetic acid 85/25/10 (v/v/v) as a liquid phase. Iodine staining was used for band visualization. The used standards were 2-dioleoyl-sn-glycero-3-phosphocholine (DOPC) 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) 1,2-dioleoyl-sn-glycero-3-phospho-L-serine (sodium salt) (DOPS); L- α -phosphatidylinositol from liver (PI) and 1,1',2,2'-tetraoleoyl cardiolipin (sodium salt) (CL) (all from Avanti Polar Lipids). Imaging of TLC plates was performed using a Luminescent Image Analyzer LAS-1000 plus (Fujifilm).

The CytcO activity was monitored by measuring the oxygen consumption using an Oxygraph Plus System (Hansatech) at 25 °C with 10 mM ascorbate, 100 μ M TMPD and 50 μ M cytochrome *c* (from equine heart, Sigma). The reaction medium contained 10 mM KCl, 5 mM MgCl₂, 10 mM KH₂PO₄ (pH 7.4), 0.1 mM EDTA and 0.3 M mannitol.

4.5. Flash photolysis and flow-flash measurements

Time-resolved kinetics experiments were performed as described e.g. in [33,54], using an LKS80 (Applied Photophysics) flash photolysis/flow-flash set-up. Briefly, the sample was transferred to an anaerobic Thunberg cuvette (path length 1.00 cm), the atmosphere was exchanged for N₂ on a vacuum line after which the CytcO in the SMA native nanodiscs was reduced with 2 mM ascorbate and 3 μ M PMS (4 mM and 1 μ M, respectively, when using mitochondria). After the reduction was complete (based on the optical absorption spectrum), the atmosphere was exchanged for carbon monoxide. To monitor CO dissociation and recombination the sample was illuminated by a laser flash (Nd-YAG laser, pulse length 8 ns, 40 mJ at 532 nm) and absorbance changes were monitored. For the flow-flash measurements, the mixing with the oxygen-saturated buffer was 1:2.5 for CytcO-SMA and 1:1 for other samples, which gives the concentration of oxygen after mixing of ~0.9 mM and 0.6 mM, respectively. The reaction buffers used were: 0.2 M NaCl, 20 mM HEPES (pH 8.0) for CytcO-SMA; 0.15 M KCl, 20 mM HEPES (pH 8.0), 0.05% DDM, 10% glycerol for the purified CytcO; 60 mM sorbitol, 20 mM HEPES (pH 7.4) for the yeast mitochondria and 0.1 M KCl, 20 mM HEPES (pH 8.0), 0.1 mM EDTA, 0.05% DDM for CytcO from bovine heart. The concentration of reactive enzyme was estimated from the amplitude of the time-unresolved absorbance changes induced by the laser flash at 445 nm using an absorption coefficient of 67 mM⁻¹ cm⁻¹.

4.6. Negative stain electron microscopy

Aliquots of frozen SMA-CytcO native nanodiscs at 4 mg/ml were thawed slowly on ice before preparing the negative stain EM grid. The sample was diluted to 0.01 mg/ml in buffer B (see above). Prior to sample application, 400-mesh Cu-Rh maxtaform grids (Electron Microscopy Sciences) that had been overlaid with a continuous carbon support were exposed to a 95% Ar/5% O₂ plasma for 20 s using a Gatan Solarus (Gatan) to generate a hydrophilic surface. A volume of 4 μ l of the sample was applied to the freshly cleaned grid. After 30 s of adsorption, excess protein was wicked away with a Whatman No. 1 filter paper (GE

Healthcare Life Sciences), and 2% (w/v) uranyl formate solution (4 μ l) was immediately applied to the grid. The grid was then inverted and placed on four 20 μ l droplets of uranyl formate solution for 10 s each, in succession. Excess stain was then wicked away with filter paper and the grid set to dry on the edge of a fume hood to provide airflow.

Grids were inserted into a Tecnai Spirit (FEI Co) LaB6 electron microscope operating at 120 keV, and images collected using a random defocus range of -0.5 to -1.5 μ m and an electron dose of 20 e⁻/Å² using the Legion automated image acquisition software (Suloway et al., 2005) at a nominal magnification of 52,000 \times (pixel size of 2.05 Å/pixel at the specimen level). 273 images were collected using a F416 CMOS 4 K \times camera (T-VIPS, Oslo, Norway). Image analysis was performed using the Appion image processing pipeline [55]. Particles were selected from the micrographs using a difference-of-Gaussians-based automated particle picking program [56] and the contrast transfer function (CTF) of each micrograph was estimated using CTFIND3 [57]. Using the estimated CTF, phases for each micrograph were corrected before extracting particles with a box size of 80 pixels. Particles were binned by a factor of two and normalized for analysis. The resulting stack of 79,343 particles was subjected to three rounds of reference-free 2D classification using iterative multivariate statistical analysis and multi-reference alignment [58], implemented in Appion. These rounds of 2D analysis were used to remove damaged, aggregated, or erroneously selected particles. The final 2D class averages result from 2D classification of the final stack of 46,763 particles. The particle size diameters of the class averages used to generate the histogram in Fig. 1 were calculated using SPIDER (PMID: 8742743). Class averages were rotationally averaged, and a 1D Radon transform was calculated in order to measure the width of the peak.

Transparency document

The Transparency document associated with this article can be found, in online version.

Acknowledgements

We would like to thank Susanna Seppälä (UCSB) for help with preparation of SMA in the initial phase of the project. This study was supported by a grant from the Knut and Alice Wallenberg Foundation (KAW 2013.0006) (to PB, CvB, PÅ, MH). Support was also obtained from the Pew Scholars program, the Searle Scholars program, and NIH grant DP2 EB020402-01 (to GCL), the Swedish Research Council (to PB, CvB, PÅ, MH), the Swedish Cancer Society (MH) and the Swedish Foundation for Strategic Research (MH).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bbame.2016.09.004>.

References

- [1] M. Zoonens, J.L. Popot, Amphipols for each season, *J. Membr. Biol.* 247 (2014) 759–796.
- [2] J.L. Rigaud, D. Lévy, Reconstitution of membrane proteins into liposomes, *METHODS IN ENZYMOLOGY*, 372, 2003, pp. 65–86.
- [3] T.K. Ritchie, Y.V. Grinkova, T.H. Bayburt, I.G. Denisov, J.K. Zolnerciks, W.M. Atkins, S.G. Sligar, Chapter 11 reconstitution of membrane proteins in phospholipid bilayer nanodiscs, *Methods in Enzymology*, 464, 2009, pp. 211–231.
- [4] S. Scheidelaar, M.C. Koorengel, J.D. Pardo, J.D. Meeldijk, E. Breukink, J.A. Killian, Molecular model for the solubilization of membranes into nanodiscs by styrene maleic acid copolymers, *Biophys. J.* 108 (2015) 279–290.
- [5] J.M. Dörr, S. Scheidelaar, M.C. Koorengel, J.J. Dominguez, M. Schäfer, C.A. van Walree, J.A. Killian, The styrene-maleic acid copolymer: a versatile tool in membrane research, *Eur. Biophys. J.* 45 (2016) 3–21.
- [6] S.C. Lee, N.L. Pollock, Membrane proteins: is the future disc shaped? *Biochem. Soc. Trans.* 44 (2016) 1011–1018.
- [7] C. Logez, M. Damian, C. Legros, C. Dupré, M. Guéry, S. Mary, R. Wagner, C. M'Kadmi, O. Nosjean, B. Fould, J. Marie, J.A. Fehrentz, J. Martinez, G. Ferry, J.A. Boutin, J.L.

- Baneires, Detergent-free isolation of functional G protein-coupled receptors into nanometric lipid particles, *Biochemistry* 55 (2016) 38–48.
- [8] A.R. Long, C.C. O'Brien, K. Malhotra, C.T. Schwall, A.D. Albert, A. Watts, N.N. Alder, A detergent-free strategy for the reconstitution of active enzyme complexes from native biological membranes into nanoscale discs, *BMC Biotechnol.* 13 (2013).
- [9] J.M. Dörr, M.C. Koorengel, M. Schäfer, A.V. Prokofyev, S. Scheidelaar, E.A.W. Van Der Crujssen, T.R. Dafforn, M. Baldus, J.A. Killian, Detergent-free isolation, characterization, and functional reconstitution of a tetrameric K⁺ channel: the power of native nanodiscs, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) 18607–18612.
- [10] T.J. Knowles, R. Finka, C. Smith, Y.P. Lin, T. Dafforn, M. Overduin, Membrane proteins solubilized intact in lipid containing nanoparticles bounded by styrene maleic acid copolymer, *J. Am. Chem. Soc.* 131 (2009) 7484–7485.
- [11] M. Jamshad, J. Charlton, Y.P. Lin, S.J. Routledge, Z. Bawa, T.J. Knowles, M. Overduin, N. Dekker, T.R. Dafforn, R.M. Bill, D.R. Poyner, M. Wheatley, G-protein coupled receptor solubilization and purification for biophysical analysis and functional studies, in the total absence of detergent, *Biosci. Rep.* 35 (2015).
- [12] M. Jamshad, V. Grimard, I. Idini, T.J. Knowles, M.R. Dowle, N. Schofield, P. Sridhar, Y. Lin, R. Finka, M. Wheatley, O.R.T. Thomas, R.E. Palmer, M. Overduin, C. Govaerts, J.M. Ruysschaert, K.J. Edler, T.R. Dafforn, Structural analysis of a nanoparticle containing a lipid bilayer used for detergent-free extraction of membrane proteins, *Nano Res.* 8 (2015) 774–789.
- [13] M. Jamshad, Y.P. Lin, T.J. Knowles, R.A. Parslow, C. Harris, M. Wheatley, D.R. Poyner, R.M. Bill, O.R.T. Thomas, M. Overduin, T.R. Dafforn, Surfactant-free purification of membrane proteins with intact native membrane environment, *Biochem. Soc. Trans.* 39 (2011) 813–818.
- [14] S. Gulati, M. Jamshad, T.J. Knowles, K.A. Morrison, R. Downing, N. Cant, R. Collins, J.B. Koenderink, R.C. Ford, M. Overduin, I.D. Kerr, T.R. Dafforn, A.J. Rothnie, Detergent-free purification of ABC (ATP-binding-cassette) transporters, *Biochem. J.* 461 (2014) 269–278.
- [15] M.C. Orwick, P.J. Judge, J. Procek, L. Lindholm, A. Graziadei, A. Engel, G. Gröbner, A. Watts, Detergent-free formation and physicochemical characterization of nanosized lipid-polymer complexes: lipodisq, *Angew. Chem. Int. Ed.* 51 (2012) 4653–4657.
- [16] M. Orwick-Rydmark, J.E. Lovett, A. Graziadei, L. Lindholm, M.R. Hicks, A. Watts, Detergent-free incorporation of a seven-transmembrane receptor protein into nanosized bilayer lipodisq particles for functional and biophysical studies, *Nano Lett.* 12 (2012) 4687–4692.
- [17] K. Skaar, H.J. Korza, M. Tarry, P. Sekyrova, M. Högbom, Expression and subcellular distribution of GFP-tagged human tetraspanin proteins in *Saccharomyces cerevisiae*, *PLoS One* 10 (2015).
- [18] D.J.K. Swainsbury, S. Scheidelaar, R. Van Grondelle, J.A. Killian, M.R. Jones, Bacterial reaction centers purified with styrene maleic acid copolymer retain native membrane functional properties and display enhanced stability, *Angew. Chem. Int. Ed.* 53 (2014) 11803–11807.
- [19] V. Postis, S. Rawson, J.K. Mitchell, S.C. Lee, R.A. Parslow, T.R. Dafforn, S.A. Baldwin, S.P. Muench, The use of SMALPs as a novel membrane protein scaffold for structure study by negative stain electron microscopy, *Biochim. Biophys. Acta Biomembr.* 1848 (2015) 496–501.
- [20] P. Brzezinski, R.B. Gennis, Cytochrome c oxidase: exciting progress and remaining mysteries, *J. Bioenerg. Biomembr.* 40 (2008) 521–531.
- [21] P. Brzezinski, A.L. Johansson, Variable proton-pumping stoichiometry in structural variants of cytochrome c oxidase, *Biochim. Biophys. Acta Bioenerg.* 1797 (2010) 710–723.
- [22] L. Näsvik Öjemyr, A. Maréchal, H. Vestin, B. Meunier, P.R. Rich, P. Brzezinski, Reaction of wild-type and Glu243Asp variant yeast cytochrome c oxidase with O₂, *Biochim. Biophys. Acta Bioenerg.* 1837 (2014) 1012–1018.
- [23] M. Vukotic, S. Oeljeklaus, S. Wiese, F.N. Vögtle, C. Meisinger, H.E. Meyer, A. Ziesenis, D.M. Katschinski, D.C. Jans, S. Jakobs, B. Warscheid, P. Rehling, M. Deckers, Rcf1 mediates cytochrome oxidase assembly and respirasome formation, revealing heterogeneity of the enzyme complex, *Cell Metab.* 15 (2012) 336–347.
- [24] V. Strogolova, A. Furness, M.R. Micaela, J. Garlich, R.A. Stuart, Rcf1 and Rcf2, members of the hypoxia-induced gene 1 protein family, are critical components of the mitochondrial cytochrome bc₁-cytochrome c oxidase supercomplex, *Mol. Cell. Biol.* 32 (2012) 1363–1373.
- [25] Y.C. Chen, E.B. Taylor, N. Dephore, J.M. Heo, A. Tonhato, I. Papandreou, N. Nath, N.C. Denko, S.P. Gygi, J. Rutter, Identification of a protein mediating respiratory supercomplex stability, *Cell Metab.* 15 (2012) 348–360.
- [26] C. Rydström Lundin, C. Von Ballmoos, M. Ott, P. Ådelroth, P. Brzezinski, Regulatory role of the respiratory supercomplex factors in *Saccharomyces cerevisiae*, *Proc. Natl. Acad. Sci. U. S. A.* 113 (2016) E4476–E4485.
- [27] R.O. Poyton, G. Schatz, Cytochrome c oxidase from bakers' yeast. III. Physical characterization of isolated subunits and chemical evidence for two different classes of polypeptides, *J. Biol. Chem.* 250 (1975) 752–761.
- [28] A. Maréchal, B. Meunier, D. Lee, C. Orengo, P.R. Rich, Yeast cytochrome c oxidase: a model system to study mitochondrial forms of the haem-copper oxidase superfamily, *Biochim. Biophys. Acta Bioenerg.* 1817 (2012) 620–628.
- [29] N. Yano, K. Muramoto, M. Mochizuki, K. Shinzawa-Itoh, E. Yamashita, S. Yoshikawa, T. Tsukihara, X-ray structure of cyanide-bound bovine heart cytochrome c oxidase in the fully oxidized state at 2.0 Å resolution, *Acta Crystallogr. F Struct. Biol. Commun.* 71 (2015) 726–730.
- [30] J.F. Nagle, S. Tristram-Nagle, Structure of lipid bilayers, *Biochim. Biophys. Acta Rev. Biomembr.* 1469 (2000) 159–195.
- [31] W.H. Vanneste, The stoichiometry and absorption spectra of components a and a-3 in cytochrome c oxidase, *Biochemistry* 5 (1966) 838–848.
- [32] B. Kadenbach, M. Hüttemann, The subunit composition and function of mammalian cytochrome c oxidase, *Mitochondrion* 24 (2015) 64–76.
- [33] C. Von Ballmoos, N. Gonska, P. Lachmann, R.B. Gennis, P. Ådelroth, P. Brzezinski, Mutation of a single residue in the ba₃ oxidase specifically impairs protonation of the pump site, *Proc. Natl. Acad. Sci. U. S. A.* 112 (2015) 3397–3402.
- [34] P. Ådelroth, P. Brzezinski, B.G. Malmström, Internal electron transfer in cytochrome c oxidase from *Rhodobacter sphaeroides*, *Biochemistry* 34 (1995) 2844–2849.
- [35] W.H. Woodruff, Coordination dynamics of heme-copper oxidases. The ligand shuttle and the control and coupling of electron transfer and proton translocation, *J. Bioenerg. Biomembr.* 25 (1993) 177–188.
- [36] H. Witt, F. Malatesta, F. Nicoletti, M. Brunori, B. Ludwig, Cytochrome-c-binding site on cytochrome oxidase in *Paracoccus denitrificans*, *Eur. J. Biochem.* 251 (1998) 367–373.
- [37] P. Ådelroth, M. Ek, P. Brzezinski, Factors determining electron-transfer rates in cytochrome c oxidase: investigation of the oxygen reaction in the *R. sphaeroides* and bovine enzymes, *Biochim. Biophys. Acta* 1367 (1998) 107–117.
- [38] M. Karpefors, P. Ådelroth, Y. Zhen, S. Ferguson-Miller, P. Brzezinski, Proton uptake controls electron transfer in cytochrome c oxidase, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 13606–13611.
- [39] M. Karpefors, P. Ådelroth, A. Aagaard, H. Sigurdson, M. Svensson Ek, P. Brzezinski, Electron-proton interactions in terminal oxidases, *Biochim. Biophys. Acta* 1365 (1998) 159–169.
- [40] C.M. Cruciat, S. Brunner, F. Baumann, W. Neupert, R.A. Stuart, The cytochrome bc₁ and cytochrome c oxidase complexes associate to form a single supercomplex in yeast mitochondria, *J. Biol. Chem.* 275 (2000) 18093–18098.
- [41] H. Schägger, Respiratory chain supercomplexes, *IUBMB Life* 52 (2002) 119–128.
- [42] R.A. Stuart, Supercomplex organization of the oxidative phosphorylation enzymes in yeast mitochondria, *J. Bioenerg. Biomembr.* 40 (2008) 411–417.
- [43] Y. Chaban, E.J. Boekema, N.V. Dudkina, Structures of mitochondrial oxidative phosphorylation supercomplexes and mechanisms for their stabilisation, *Biochim. Biophys. Acta Bioenerg.* 1837 (2014) 418–426.
- [44] J. Heinemeyer, H.P. Braun, E.J. Boekema, R. Kouřil, A structural model of the cytochrome c reductase/oxidase supercomplex from yeast mitochondria, *J. Biol. Chem.* 282 (2007) 12240–12248.
- [45] D.R. Winge, Sealing the mitochondrial respirasome, *Mol. Cell. Biol.* 32 (2012) 2647–2652.
- [46] E. Mileykovskaya, P.A. Penczek, J. Fang, V.K.P.S. Mallampalli, G.C. Sparagna, W. Dowhan, Arrangement of the respiratory chain complexes in *Saccharomyces cerevisiae* supercomplex III₂IV₂ revealed by single particle cryo-electron microscopy, *J. Biol. Chem.* 287 (2012) 23095–23103.
- [47] M.L. Genova, G. Lenaz, Functional role of mitochondrial respiratory supercomplexes, *Biochim. Biophys. Acta Bioenerg.* 1837 (2014) 427–443.
- [48] R. Acín-Perez, J.A. Enriquez, The function of the respiratory supercomplexes: the plasticity model, *Biochim. Biophys. Acta Bioenerg.* 1837 (2014) 444–450.
- [49] H. Boumans, L.A. Grivell, J.A. Berden, The respiratory chain in yeast behaves as a single functional unit, *J. Biol. Chem.* 273 (1998) 4872–4877.
- [50] C. Meisinger, N. Pfanner, K.N. Truscott, Isolation of yeast mitochondria, *Methods Mol. Biol.* 313 (2006) 33–39.
- [51] M. Björck, S. Zhou, C. Rydström Lundin, M. Ott, P. Ådelroth, P. Brzezinski, Kinetics of CO binding to flavohemoglobin in different cellular compartments of *S. cerevisiae* (submitted for publication) 2016.
- [52] B. Meunier, A. Maréchal, P.R. Rich, Construction of histidine-tagged yeast mitochondrial cytochrome c oxidase for facile purification of mutant forms, *Biochem. J.* 444 (2012) 199–204.
- [53] U. Brandt, H. Schägger, G. von Jagow, Purification of cytochrome-c oxidase retaining its pulsed form, *Eur. J. Biochem.* 182 (1989) 705–711.
- [54] J. Vilhjálmsson, A.L. Johansson, P. Brzezinski, Structural changes and proton transfer in cytochrome c oxidase, *Sci. Rep.* 5 (2015).
- [55] G.C. Lander, S.M. Stagg, N.R. Voss, A. Cheng, D. Fellmann, J. Pulokas, C. Yoshioka, C. Irving, A. Mulder, P.W. Lau, D. Lyumkis, C.S. Potter, B. Carragher, Appion: an integrated, database-driven pipeline to facilitate EM image processing, *J. Struct. Biol.* 166 (2009) 95–102.
- [56] N.R. Voss, C.K. Yoshioka, M. Radermacher, C.S. Potter, B. Carragher, DoG picker and TiltPicker: software tools to facilitate particle selection in single particle electron microscopy, *J. Struct. Biol.* 166 (2009) 205–213.
- [57] J.A. Mindell, N. Grigorieff, Accurate determination of local defocus and specimen tilt in electron microscopy, *J. Struct. Biol.* 142 (2003) 334–347.
- [58] T. Ogura, K. Iwasaki, C. Sato, Topology representing network enables highly accurate classification of protein images taken by cryo electron-microscope without masking, *J. Struct. Biol.* 143 (2003) 185–200.
- [59] W.L. DeLano, The PyMOL Molecular Graphics System, DeLano Scientific, Palo Alto, CA, USA, 2002.